

UNCLASSIFIED

AD. 295 124

*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGED BY ASTIA
295 124
AS AD NO. _____

295 124

OFFICE OF NAVAL RESEARCH

Contract Nonr 3471(00)

Task No. 356-433

TECHNICAL REPORT NO. 9

The Effect of Substitution on the Ionization Potentials of
Free Radicals and Molecules. V Extension of the Density
Matrix Treatment to a Two-Body Model for Electrons
Outside a "Core".

by

Joyce J. Kaufman

December, 1962

RIAS
Research Institute for Advanced Study
7212 Bellona Avenue
Baltimore 12, Maryland

Reproduction in whole or in part is permitted
for any purpose of the United States Government.

Abstract

Extension has been made to a two-body model for a pair of electrons outside a "core" of the treatment by density matrix formalism of a one-body model for an electron outside a "core". δ_k values [differences in ionization potentials between the parent compound] for the π electron of alkyl radicals can be developed exactly as a one-body model problem in terms of density matrices. The derivation of the two-body model permits one quantum mechanically to describe the same δ_k effects on the ionization potentials of an electron from the lone pair of substituted amines by an analogous treatment using the same perturbation Hamiltonian as for substituted alkyl radicals with two electron functions for the lone pair of the nitrogen.

Introduction

In part II of the series¹ the constancy of δ_k values [which measure the change in ionization potential between a substituted molecule (or radical) and the parent compound] for alkyl radicals and amines was discussed in relation to the energies and possible wave functions for these species and their positive ions in terms of ordinary first order perturbation theory. In that article it was assumed that at least in the alkyl radicals one could separate the σ and π electrons and, to a first approximation, deal with the Hamiltonian, wave function, and orbital of only the π electron. The basis for the above assumption at that time was that this separation is the one commonly invoked in virtually all theoretical treatments of π systems. However, verification for the separation in the case of alkyl radicals can now be based on the recent justification of the one-body model for an electron outside a "core"² which was derived in terms of density matrices. The problem of the alkyl radicals can also be developed exactly as a one-body model in terms of density matrix formalism.

In II it was shown to be possible to describe the wave functions and Hamiltonians for the substituted and unsubstituted amines in a manner which emphasizes their close similarity to the alkyl radicals by making a formal separation into σ and " π -like" electrons, where " π -like" implied only that this system is orthogonal to the other bonds in the molecule. This division depended upon the proven fact that it is possible to write the wave function for the

complete molecule as a product of wave functions which must be mutually orthogonal for the separate bonds or lone pairs so that certain orbitals may be associated with selected parts.³ Later in this article it will be seen that the one-body model suitable for describing the alkyl radicals can be extended to a two-body model for a pair of electrons outside a "core" suitable for describing the amines.

Extension to Two-Body Model

The pertinent parts of the article on the one-body model² are summarized here to make it easier to follow its extension to the two-body model and its subsequent application to the alkyl radicals and amines.

Resume of One-Body Model

The system comprising a core and one valence electron was described by a wave function

$$\Psi(1,2,\dots,N) = M \mathcal{A} \{ \Phi_A(1,2,\dots,[N-1]) \Phi_B(N) \} \quad (1)$$

where $\Phi_A(1,2,\dots,[N-1])$ is an antisymmetrical core function, $\Phi_B(N)$ is the spin orbital of the valence electron, \mathcal{A} is the antisymmetrizer and M is the normalization factor. For convenience Φ_A and Φ_B were individually assumed to be normalized. No other assumptions were made and although (1) cannot be an exact wave function it provides a rather general description of a system with two distinguishable parts. The variational solution of

$$h_{eff} \Psi = \epsilon f \phi \quad (2)$$

proceeded even without orthogonality restrictions.

The density matrices of the whole system can be expressed in terms of those of its separate parts.

Definition of density matrix:

$$\rho_n(1,2,\dots,n;1',2',\dots,n') = \frac{N!}{(N-1)!} \int \Psi(1,\dots,N) \Psi^*(1',\dots,n';\overline{n+1},\dots,N) dx_{n+1}\dots dx_N \quad (3)$$

With analogous group density matrices ρ_n^A and ρ_n^B one obtained

$$\rho_1(1;1') = \{\rho_1^A(1;1') + \rho_1^B(1;1')\} - \text{-----} \quad (4)$$

$$\begin{aligned} \rho_2(1,2;1',2') = & \{\rho_2^A(1,2;1',2') + \rho_1^A(1;1') \rho_1^B(2;2') \\ & - \rho_1^A(2;1') \rho_1^B(1;2') + \rho_1^B(1;1') \rho_1^A(2;2') \\ & - \rho_1^B(2;1') \rho_1^A(1;2')\} - \text{-----} \end{aligned} \quad (5)$$

neglecting normalization constants and keeping only terms which remain when the functions Φ_A and Φ_B are orthogonal in the strong sense:

$$\int \Phi_A(1,i,j\cdots) \Phi_B(1) dx_1 = 0 \quad (6)$$

The remaining terms would arise from penetration of the valence electron into the core.

The energy of the whole system is given by

$$E = \text{tr } h(1) \rho_1(1;1') + \frac{1}{2} \text{tr } g(1,2) \rho_2(1,2;1',2') \quad (7)$$

where $h(1)$ is a modified one-body operator which depends on the core wave function, $g(1,2) = \frac{e^2}{r_{12}}$ and tr indicates removal of primes (after operation) and integration over all remaining variables.

It was then shown that

$$E = E^A + E_{\text{eff}}^B = E^A + \tilde{E}^B \quad (8)$$

where E^A is the energy of the core by itself

$$E^A = \text{tr } h(1) \rho_1^A(1;1') + \frac{1}{2} \text{tr } g(1,2) \rho_2^A(1,2;1',2') \quad (9)$$

and \tilde{E}^B is the energy of the valence electron itself in an effective field due to the core.

$$\begin{aligned} \tilde{E}_B = & \text{tr } h(1) \rho_1^B(1;1') + \text{tr} \int g(1,2) \rho_1^A(2;2) dx_2 \rho_1^B(1;1') \\ & - \text{tr} \int g(1,2) \rho_1^A(1;2) \rho_1^B(2;1') dx_2 \end{aligned} \quad (10)$$

\tilde{E}^B was expressed as

$$\tilde{E}_B = \text{tr } \tilde{h}^B(1) \rho_1^B(1,1') \quad (11)$$

where $\tilde{h}^B(1)$ is an integral operator defined by

$$\tilde{h}^B(1) \phi(1) = \int \tilde{h}^B(1;2) \phi(2) dx_2 \quad (12)$$

$\tilde{h}^B(1;2)$ - the kernel - consists of several terms and the operator is a corresponding sum of operators

$$\tilde{h}^B(1) = h(1) + \tilde{A}(1) - \cancel{A}(1) + \text{penetration terms} \quad (13)$$

$$\int \tilde{A}(1) \rightarrow J^A(1) = \int g(1,2) \rho_1^A(2;2) dx_2 \quad (14)$$

$$\cancel{V}^A(1) \rightarrow K^A(1;2) = g(1,2) \rho_1^A(1;2) \quad (15)$$

$\int V^A(1)$ and $\int \cancel{V}^A(1)$ are the Coulomb and exchange potentials due to the core which are present even when (6) is satisfied.

Two-Body Model

For the two-body model for a pair of electrons outside a "core" keeping only the terms which remain when ϕ_A and ϕ_B are orthogonal according to (6), one now gets an additional term in the second order density matrix (5)

$$+ \rho_2^B(1,2;1',2') \quad (16)$$

When the expressions for the first and second order density matrices are substituted into the formula for the energy the following expression is obtained for the integral operator $\tilde{h}^B(1)$

$$\tilde{h}^B(1) = h(1) + \int V^A(1) - \cancel{V}^A(1) + \frac{1}{2} \int V^B(1) - \frac{1}{2} \cancel{V}^B(1) \quad (17)$$

where it must now be remembered that there is a pair of electrons in ϕ_B instead of just one electron.

Application of One-Body Model to Alkyl Radicals

Under the assumption that $S^{AB} = 0$

$$E^B = E - E^A = \frac{(\phi_B | h_{eff}^B | \phi_B)}{(\phi_B | \phi_B)} \quad (18)$$

where $h_{eff}^B = \tilde{h}^B$.

$-E^B$ represents the ionization potential (I.P) of the lone electron in ϕ_B which in our case is the ionization potential of the single electron in the π orbital of the alkyl radical. The derivation is completely rigorous for the alkyl radicals since $S^{AB} = 0$ for these radicals.

Since ϕ_B has been already been assumed to be normalized:

$$\text{I.P.} = -E_B = \int \phi_B(1) h_{\text{eff}}^B(1) \phi_B(1) d\tau_1 \quad (19)$$

This is identical to the expression used in II for the calculation of ionization potentials of alkyl radicals where we used only the wave function of the π electron and an h effective for the π electron.

For substituted alkyl radicals one merely changes the h_{eff}^B . To describe the effects of alkyl substituents, we assumed that one could make a perturbation on h_{eff} for the parent methyl radical

$$\begin{array}{ccccc} h_{\text{eff}} & = & h_{\text{eff}} & + & h'_{\text{eff}} \\ (\text{YCH}_2) & & (\text{CH}_3) & & (\text{H} \rightarrow \text{Y}) \end{array} \quad (20)$$

and use first order perturbation theory to evaluate the elements of h'_{eff} .

Our main purpose in II was to develop a method for describing the effects of substituent groups on the ionization potentials of substituted alkyl radicals and the corresponding amines which emphasized what must be at least a close formal similarity between the two since identical substituent groups produce almost identical perturbations on the ionization potentials.

Application of Two-Body Model to Amines

Even if ϕ_A and ϕ_B are not rigorously orthogonal it was shown² that the final expression for the energy is very similar in form

$$E - E^A = \frac{(\phi_B | h_{eff}^B | \phi_B)}{(\phi_B | f^B | \phi_B)} \quad (21)$$

where now $h_{eff}^B = \tilde{h}^B + E^A \rho_1^A$

$$f^B = 1 - \rho_1^A$$

with $\rho_1^A(1)$ the density operator with kernel $\rho_1^A(1;2)$.

From the expression

$$E = E^A + \frac{\tilde{E}^B - S^{AB} E^A}{1 + S^{AB}} \quad (22)$$

it can be seen that the closer $S^{AB} \rightarrow 0$, the closer will (22) approach (19).

For substituted amines we are changing h_{eff}^B by the same perturbation as we are changing h_{eff} for alkyl radical

$$\begin{array}{ccccc} h_{eff} & = & h_{eff} & + & h'_{eff} \\ (YCH_2) & & (CH_3) & & (H \rightarrow Y) \end{array} \quad (20)$$

$$\begin{array}{ccccc} h_{eff} & = & h_{eff} & + & h'_{eff} \\ (YCH_2) & & (NH_3) & & (H \rightarrow Y) \end{array} \quad (23)$$

In each case we are evaluating only the elements of the same h'_{eff} with respect to the alkyl radical or amine wave functions respectively (to calculate the ionization potential itself of an amine E^B would be evaluated for only one of the electrons in the nitrogen lone pair.) The basic assumption was made in part II that one could use a wave function for the π electron of the methyl radical

$$\chi_{\pi}^{\circ} = |\psi_1(1) \psi_1(0)| \quad (24)$$

$\Gamma(\text{CH}_3)$

and the analogous expression

$$\chi_{\pi}^{\circ} = M |\psi_1(1) \psi_1(1')| \quad (25)$$

(NH_3)

for the lone pair electrons in amines. " π -like" implied only that the " π -like" system is orthogonal to the other bonds in the molecule.

The perturbation calculation was outlined in part II treating integrals over the one electron χ_{π}° functions and over the two (CH_3)

electron χ_{π}° functions in exactly the same formalistic manner for (NH_3)

identical perturbing groups. Intuitively this approach had to be fruitful because from experiments we know that the same perturbing group produced the same change in ionization potential (δ_k value) regardless of the core on which it was placed.

Now, by examination of the derivation of the one-body model for an electron outside a "core" and our extension to the two-body model for a pair of electrons outside a "core", it can be seen that nowhere is it invoked that ϕ_B be a one electron function. It is just as valid to have

$$\Psi'(1,2,\dots,N) = M' \left\{ \phi_A(1,\dots,[N-2]), \phi_B([N-1],N) \right\}$$

which is simply a general description of a system with two distinguishable parts. ϕ_B is analogous to χ^* whether it be χ^*_{π} (CH₃) a one electron function, or $\chi^*_{\pi\pi}$, a two electron function. (NH₃)

Integrals over the χ 's can be treated in exactly the same way. The only thing that is important is that the h^B_{eff} be defined correctly to include all the interaction terms of the second electron. For our calculations of the δ_k values of the amines all of the electron interaction terms will approximately cancel since we are looking only at differences. The most vital point is the justification for our having treated integrals over the two electron functions in the same manner as those over the one electron functions.

Bibliography

1. "The Effect of Substitution on the Ionization Potentials of Free Radicals and Molecules. II. Theoretical Interpretation of δ_k Values for Alkyl Radicals and Amines". Joyce J. Kaufman. Presented before the Symposium on the Structure and Reactivity of Small Molecular Species at the 18th International Congress of Pure and Applied Chemistry, Montreal, Canada, August, 1961. Also RIAS Technical Report TR 62-8 April 1962.
2. "Justification of the One-Body Model for An Electron Outside a "Core" with Applications to Lithium and Sodium." Y. Ohrn and R. McWeeny. Technical Note 60. Quantum Chemistry Group. Uppsola, Sweden. February 15, 1961.
3. C. A. Coulson and M. J. Stephen, Trans. Faraday. Soc., 53 272 (1957).

Acknowledgements

The author would like to thank the Office of Naval Research
and the Soroptimist Foundation for support.

TECHNICAL REPORT DISTRIBUTION LIST

Contract - Nonr 3471(00)

NR # 356-433

No. of Copies

Commanding Officer
Office of Naval Research Branch Office
The John Crerar Library Building
86 East Randolph Street
Chicago 1, Illinois 1

Commanding Officer
Office of Naval Research Branch Office
346 Broadway
New York 13, New York 1

Commanding Officer
Office of Naval Research Branch Office
1030 East Green Street
Pasadena 1, California 1

Commanding Officer
Office of Naval Research Branch Office
Box 39 Navy # 100 Fleet Post Office
New York, New York 7

Director, Naval Research Laboratory
Washington 25, D. C.

Attn: Technical Information Officer 6
Chemistry Division 2

Chief of Naval Research
Department of the Navy
Washington 25, D. C.
Attn: Code 425 2

DDR&E
Technical Library
Room 3C-128, The Pentagon
Washington 25, D. C. 1

Technical Director
Research & Engineering Division
Office of the Quartermaster General
Department of the Army
Washington 25, D. C. 1

Technical Report Distribution List ... cont.Contract Nonr 3471(00)

	<u>No. of Copies</u>
Research Director Clothing & Organic Materials Division Quartermaster Research & Engineering Command U. S. Army Natick, Massachusetts	1
Air Force Office of Scientific Research (SRC-E) Washington 25, D. C.	1
Commanding Officer Diamond Ordnance Fuze Laboratories Washington 25, D. C. Attn: Technical Information Office Branch 012	1
Office, Chief of Research & Development Department of the Army Washington 25, D. C. Attn: Physical Sciences Division	1
Chief, Bureau of Ships Department of the Navy Washington 25, D. C. Attn: Code 342C	2
Chief, Bureau of Naval Weapons Department of the Navy Washington 25, D. C. Attn: Technical Library Code RRMA-3	3 1
ASTIA Document Service Center Arlington Hall Station Arlington 12, Virginia	10
Director of Research U.S. Army Signal Research & Development Laboratory Fort Monmouth, New Jersey	1
Naval Radiological Defense Laboratory San Francisco 24, California Attn: Technical Library	1

Technical Report Distribution List ... cont.Contract Nonr 3471(00)No. of Copies

Naval Ordnance Test Station	
China Lake, California	
Attn: Head, Chemistry Division	1
Code 40	1
Code 50	1
Commanding Officer	
Army Research Office	
Box CM, Duke Station	
Durham, North Carolina	
Attn: Scientific Synthesis Office	1
Brookhaven National Laboratory	
Chemistry Department	
Upton, New York	1
Atomic Energy Commission	
Division of Research	
Chemistry Programs	
Washington 25, D. C.	1
Atomic Energy Commission	
Division of Technical Information Extension	
Post Office Box 62	
Oak Ridge, Tennessee	1
U. S. Army Chemical Research and Development Labs.	
Technical Library	
Army Chemical Center, Maryland	1
Office of Technical Services	
Department of Commerce	
Washington 25, D. C.	1
Bureau of Naval Weapons Representative	
Westinghouse Electric Corporation	
P. O. Box 746	
Baltimore 3, Maryland	1
Commanding Officer	
Office of Naval Research Branch Office	
1000 Geary Street	
San Francisco 9, California	
Attn: Dr. P. A. Miller	1

Technical Report Distribution List ... cont.Contract Nonr 3471(00)No. of Copies

Commanding Officer
Office of Naval Research Branch Office
495 Summer Street
Boston 10, Massachusetts
Attn: Dr. A. L. Powell

1

Dr. H. S. Gutowsky
Department of Chemistry
University of Illinois
Urbana, Illinois

1

Dr. D. A. Brown
Department of Chemistry
University College
Upper Merrion Street
Dublin, Ireland

1

Dr. J. T. Vanderslice
Institute of Molecular Physics
University of Maryland
College Park, Maryland

1

Dr. Anton B. Burg
Department of Chemistry
University of Southern California
University Park
Los Angeles 7, California

1

Dr. E. G. Rochow
Department of Chemistry
Harvard University
Cambridge 38, Massachusetts

1

Dr. T. D. Parsons
Department of Chemistry
Oregon State College
Corvallis, Oregon

1

Dr. T. P. Onak
Department of Chemistry
Los Angeles State College
Los Angeles, California

1

Technical Report Distribution List ... cont.Contract Nonr 3471(00)No. of Copies

Dr. Robert W. Taft, Jr. Department of Chemistry Pennsylvania State University University Park, Pennsylvania	1
Dr. J. D. Roberts Department of Chemistry California Institute of Technology Pasadena, California	1
Dr. J. E. Leffler Department of Chemistry Florida State University Tallahassee, Florida	1
Dr. M. J. S. Dewar Department of Chemistry University of Chicago Chicago, Illinois	1
Dr. T. L. Heying Olin Mathieson Chemical Corporation 275 Winchester Avenue New Haven, Connecticut	1
Dr. G. Barth-Wehrenalp, Director Inorganic Research Department Pennsalt Chemicals Corporation Box 4388 Philadelphia 18, Pennsylvania	2
Mr. K. Warren Easley Monsanto Research Corporation Everett Station Boston 49, Massachusetts	1
Dr. Hans H. Jaffe Department of Chemistry University of Cincinnati Cincinnati, Ohio	1
Dr. Murray S. Cohen Chief, Propellant Synthesis Section Reaction Motors Division Thiokol Chemical Corporation Denville, New Jersey	1

Technical Report Distribution List ... cont.Contract Nonr 3471(00)No. of Copies

Professor W. S. Koski
Chemistry Department
The Johns Hopkins University
Baltimore, Maryland

1

Dr. Vernon H. Dibeler, Chief
Mass Spectrometry Section
National Bureau of Standards
Washington 25, D. C.

1

Dr. E. J. Prosen, Chief
Thermochemistry Section
National Bureau of Standards
Washington 25, D. C.

1

Dr. L. J. Edwards
Manager of Programs
Callery Chemical Company
Callery, Pennsylvania

1

Dr. Edgar W. Fajans
Assoc. Director of Chemical Research
U.S. Borax Research Corporation
412 Crescent Way
Anaheim, California

1

Dr. R. W. Parry
Department of Chemistry
University of Michigan
Ann Arbor, Michigan

1

Commanding Officer
Frankford Arsenal
Philadelphia 37, Pennsylvania

Attn: ORDBA-LCO

1

Dr. Riley Schaeffer
Department of Chemistry
Indiana University
Bloomington, Indiana

1

Dr. S. Golden
Department of Chemistry
Brandeis University
Waltham 54, Massachusetts

1

Technical Report Distribution List ... cont.Contract Nonr 3471(00)No. of Copies

Dr. M. F. Hawthorne Rohm and Haas Company Philadelphia 5, Pennsylvania	1
Dr. W. N. Lipscomb Department of Chemistry Harvard University Cambridge 38, Massachusetts	1
Dr. H. C. Clark Department of Chemistry University of British Columbia Vancouver, British Columbia Canada	1
Dr. H. B. Jonassen Department of Chemistry Tulane University New Orleans, Indiana	1
Dr. M. M. Chamberlain Department of Chemistry Western Reserve University Cleveland, Ohio	1
Dr. D. C. Bradley Department of Chemistry University of Western Ontario London, Canada	1
Dr. Henry Taube Department of Chemistry Stanford University Stanford, California	1
Dr. C. E. Nordmann Department of Chemistry University of Michigan Ann Arbor, Michigan	1
Dr. Ross I. Wagner American Potash & Chem. Corporation Research Department Whittier, California	1

Technical Report Distribution List ... cont.Contract Nonr 3471(00)No. of Copies

Dr. E. C. Ashby
Ethyl Corporation
Baton Rouge, Louisiana

1

Dr. George F. Huff, Vice President
Research and Development
Callery Chemical Company
Callery, Pennsylvania

1

Dr. David Chamberlain, Jr.
Stanford Research Institute
Menlo Park, California

1